

Green LEP vs ThuLEP in Management of Marked Enlarged Prostate

NCT03305861

Status	RECRUITING
Phase	Not Applicable
Sponsor	Mansoura University
Enrollment	110 participants

Key Eligibility Criteria

Inclusion (6)

- Patients' age \geq 40 years
- LUTS secondary to BOO due to BPH who failed medical treatment
- International prostate symptom scores (IPSS) \geq 15 and bother score (QOL) \geq 3 (according to IPSS question 8)
- Peak urinary flow rate (Qmax) $<$ 15 ml/sec with at least 125 ml voided volume or Patients with acute urine retention secondary to BPH who failed trial of voiding on medical treatment.
- ASA (American society of anaesthesiologists) score \leq 3.

... and 1 more (see full listing online)

Exclusion (7)

- Patient with neurological disorder which might affect bladder function as cerebrovascular stroke, Parkinson disease
- Active urinary tract infection,
- Presence of active bladder cancer (within the last 2 years)
- Known prostate cancer patients will be excluded preoperatively on the basis of digital rectal examination, prostate specific antigen level, and TRUS imaging followed by prostate biopsies if necessary.
- Patient has a disorder of the coagulation cascade (e.g., liver cell failure) or disorders that affect platelet count or function (e.g., von Willebrand disease) that would put the subject at risk for intraoperative or postoperative bleeding.

... and 2 more (see full listing online)

Locations (1 total)

Urology and Nephrology Center, Al Mansurah, Dakahlia Governorate, Egypt