

Effects of Sucralose on Drug Absorption and Metabolism (The Sweet-Meds Study)

NCT03407079

Status	RECRUITING
Phase	Phase 2
Sponsor	National Institute on Alcohol Abuse and Alcoholism (NIAAA)
Enrollment	150 participants

Key Eligibility Criteria

Inclusion (11)

- Age: between 18 and 60 years
 - Female adults who self-identify as Hispanic and/or Black
 - Body weight greater than 50 kg (110 lb)
 - Body mass index between 25 kg/m² and 40 kg/m²
 - Consumption of less than or equal to one 12-ounce beverage sweetened with NNS per month or food equivalent
- ... and 6 more (see full listing online)

Exclusion (13)

- Current use of prescription or non-prescription medication(s), herbal medications and oral contraceptives are also excluded. Certain exceptions are permitted, including vitamins. Other medications may be permitted at the discretion of the investigators.
 - Diabetes (fasting blood glucose of 126 mg/dl or higher, or 2-hour blood glucose of 200 or higher on OGTT)
 - Taken medications that affect blood sugar in the past 3 months or that include antibiotics
 - GI history, at the discretion of the investigators
 - Known allergy, sensitivity, or other contraindication to study procedures
- ... and 8 more (see full listing online)

Locations (1 total)

National Institutes of Health Clinical Center, Bethesda, Maryland, United States