

# Low- and Moderate-intensity Statin and Clinical Outcome of Primary Prevention in Individuals Aged >70 Years: the SCOPE-70 RCT Study

NCT03770312

---

Status	RECRUITING
Phase	Phase 4
Sponsor	Yonsei University
Enrollment	724 participants

## Key Eligibility Criteria

---

### Inclusion (12)

- Adults aged 70 years or older as of the date of written consent, meeting one of the criteria in items 2-5)
  - Patients with no prior statin use or who have discontinued statin therapy for at least 4 weeks, with no cardiovascular risk factors\* and LDL-cholesterol levels of 160-189 mg/dL
  - Patients with no prior statin use or who have discontinued statins for at least 4 weeks, with one or more cardiovascular disease risk factors\* and LDL-cholesterol levels of 80-189 mg/dL
  - For patients currently taking statins, with 0 cardiovascular disease risk factors\* and LDL-cholesterol levels of 95-114 mg/dL
  - For patients currently taking statins, with 1 or more cardiovascular disease risk factors\* and LDL-cholesterol levels of 50-114 mg/dL
- ... and 7 more (see full listing online)

### Exclusion (9)

- Individuals who have been diagnosed with cancer within the last 5 years.
  - Individuals who have high level of serum ALT ( $\geq 2$  upper normal limit).
  - Individuals who have serum creatinine  $\geq 2$  mg/dL.
  - Individuals who have been diagnosed with cardiovascular disease ( $\geq 50\%$  stenosis of coronary artery on imaging study, percutaneous transluminal coronary angioplasty, coronary artery bypass graft, or myocardial infarction).
  - Individuals who have been diagnosed with peripheral artery disease ( $\geq 50\%$  stenosis of peripheral artery on imaging study, or ankle brachial index  $\leq 0.9$  or  $\geq 1.3$ ).
- ... and 4 more (see full listing online)

## Locations (1 total)

---

Division of Cardiology, Cardiovascular Center, Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea