

# Skeletal Muscle Function in Interstitial Lung Disease

NCT03800017

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	University of British Columbia
<b>Enrollment</b>	40 participants

## Key Eligibility Criteria

---

### Inclusion (12)

- Age 40-80 years (inclusive)
- A multidisciplinary diagnosis of idiopathic pulmonary fibrosis (IPF), idiopathic fibrotic nonspecific interstitial pneumonia (NSIP), chronic hypersensitivity pneumonitis (HP), or unclassifiable ILD with a differential diagnosis that consists of the above diagnoses
- Fibrosis on high resolution computed tomography (HRCT): honeycombing, reticulation, or traction bronchiectasis
- Appropriate candidate for pulmonary rehabilitation
- minute walk distance 50m or more
- ... and 7 more (see full listing online)

### Exclusion (9)

- Contraindication to exercise testing (e.g. significant cardiovascular, musculoskeletal, neurological disease)
- Other significant extra-pulmonary disease that, based on clinical assessment, could impair exercise capacity and/or oxygenation
- Forced vital capacity (FVC) less than 50% or Diffusion capacity for carbon monoxide (DLCO) less than 25%
- Concurrent or recent participation (less than 6 months) in a pulmonary rehabilitation program
- Use of prednisone greater than 10 mg/day for more than 2 weeks within 3 months of the first study visit
- ... and 4 more (see full listing online)

## Locations (1 total)

---

St. Paul's Hospital, Vancouver, British Columbia, Canada