

Comparing the Efficacy of Steroid, Acupuncture and Platelet Rich Plasma Injection in Patients with Sciatica

NCT03801512

Status	RECRUITING
Phase	Not Applicable
Sponsor	Taipei Medical University Hospital
Enrollment	90 participants

Key Eligibility Criteria

Inclusion (6)

- Female or male, 20 to 80 years of age.
- Participants who had dermatome between L3 to S1, or participants whose straight leg raising test was positive, or participants had Meyerding Grade I to II spondylolisthesis or herniated intervertebral disc or spinal stenosis diagnosed by X-rays, Magnetic resonance imaging or computed tomography.
- Participants whose sciatica symptom doesn't have occur time limit.
- Participants who agreed to follow the trial protocol.
- Participants who could complete the study treatment and assessments.

... and 1 more (see full listing online)

Exclusion (6)

- Participants with severe progressive neurological symptoms (e.g. cauda equina compression).
- Participants who had undergone surgery for lumbar disc herniation within 6 months.
- Participants whose sciatica symptom wasn't caused by lumbar radiculopathy.
- Participants with cardiovascular, liver, kidney, or hematopoietic system diseases, severe psychiatric or psychological disorder, or cancer.
- Women who were pregnant.

... and 1 more (see full listing online)

Locations (1 total)

Taipei Medical University Hospital, Taipei, Taiwan, Taiwan