

# Effects of Testosterone Undecanoate vs Placebo on Intrahepatic Fat Content in Overweight/Obese Men With T2DM or Prediabetes and Hypogonadism

NCT03851627

---

Status	RECRUITING
Phase	Phase 4
Sponsor	Alexandra Kautzky-Willer
Enrollment	32 participants

## Key Eligibility Criteria

---

### Inclusion (8)

- prediabetes/T2DM
- male sex
- HbA1c  $\geq 5.7\%$  -  $9.0\%$  or fasting glucose  $\geq 100\text{mg/dl}$  or postprandial glucose  $\geq 140\text{mg/dl}$
- Age  $\geq 18$  -  $75$  years
- BMI  $\geq 25\text{kg/m}^2$
- ... and 3 more (see full listing online)

### Exclusion (32)

- Current testosterone treatment or testosterone replacement within the last 12 month
- Serum creatinine  $> 1.5\text{mg/dl}$
- Liver enzymes above 3 fold normal range
- PSA  $> 4.0\text{ng/ml}$
- Hematocrit  $> 50\%$
- ... and 27 more (see full listing online)

## Locations (1 total)

---

Abt. für Endokrinologie & Stoffwechsel, Univ. Klin f. Innere Medizin III, Vienna, Austria