

Von Willebrand Factor in Pregnancy (VIP) Study

NCT04146376

Status	RECRUITING
Sponsor	University of Washington
Enrollment	110 participants

Key Eligibility Criteria

Inclusion (6)

- von Willebrand Disease (VWD) patients defined prepartum as Type 1 per National Heart, Lung, and Blood Institute (NHLBI) criterion of von Willebrand Factor (VWF) level less than 30 percent, or Type 2, or Type 3 VWD
 - or
 - A diagnosis of VWD and VWF and Factor VIII (FVIII) levels obtained in gestational weeks 34-38 that determine enrollment in the non-corrector group:
 - Patients with gestational week 34-38 VWF:Ag, VWF:Act (or VWF:RCO), or FVIII:C less than 100 percent will be enrolled in the non-corrector group. In patients with an isolated VWF:CB type 2 defect, VWF:CB less than 100 percent can also be determined as a non-corrector
 - Patients with all VWF parameter levels greater than or equal to 100 percent self-corrected at gestational weeks 34-38 will be enrolled in the corrector group
- ... and 1 more (see full listing online)

Exclusion (5)

- Presence of a clinical contraindication to receive wilate or tranexamic acid, as determined by the health care provider, such as a prior drug reaction
- Presence of other concurrent disorder of hemostasis, platelet dysfunction, or collagen disorders
- Presence of liver disease or renal disease, clinical suspicion or diagnosis of preeclampsia or eclampsia, HELLP syndrome, TTP, DIC, or other acquired vasculopathy or coagulopathy
- Age less than 18 years
- Inability of the local laboratory to monitor the VWF laboratory tests needed during the course of treatment to determine Wilate dosing adjustments

Locations (11 total)

University of Colorado, Aurora, Colorado, United States
Yale University, New Haven, Connecticut, United States
University of Miami, Miami, Florida, United States
... and 8 more locations