

Fragility Fractures of the Pelvis (FFP)

NCT04182776

Status	RECRUITING
Sponsor	AO Innovation Translation Center
Enrollment	420 participants

Key Eligibility Criteria

Inclusion (7)

- Age 65 years or older at time of injury
- Diagnosis, via CT scan as per standard of care, of FFP grade II or higher according to the Rommens and Hofmann classification (including fractures of the anterior pelvic ring involving the anterior lip of the acetabulum not regarded as the acetabular fractures)
- Informed consent obtained, ie:
- Ability to understand the content of the patient information/ICF
- Willingness and ability to participate in the registry according to the standard of care in each clinic
- ... and 2 more (see full listing online)

Exclusion (6)

- Pelvic fractures that do not meet the criteria of a fragility fracture (ie, high-energy trauma)
- FFP type I fractures
- Concomitant fractures of the acetabulum (except anterior lip of the acetabulum, see above)
- Pathological fractures (ie, patients with malignant or metastatic diseases of the pelvis, infections)
- Concomitant osteoporotic fractures outside of pelvis
- ... and 1 more (see full listing online)

Locations (10 total)

University of Missouri, Columbia, Missouri, United States
The Ottawa Hospital - Civic Campus, Ottawa, Canada
University Medical Center Hamburg Eppendorf, Hamburg, Germany
... and 7 more locations