

# Effect of Giving Reduced Fluid in Children After Trauma

NCT04201704

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Columbia University
<b>Enrollment</b>	250 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- Trauma patients older than 6 months and younger than 15 years admitted to the pediatric intensive care unit (PICU)
- Patients admitted to the PICU directly from the Emergency Department (ED)
- Patients admitted to the PICU from the operating room (OR)
- Patients transferred to PICU from outside facility ED (need to have been in ED 12 hours or less)

### Exclusion (11)

- Patients transferred to PICU from outside PICU or inpatient floor
- Patients transferred to PICU from outside facility ED if >12 hours
- Patients expected to be discharged from the PICU within 24 hours
- Patient with congenital heart disease as defined by a congenital cardiac defect requiring surgery or medication
- Patient with diagnosis of chronic cardiac condition (e.g. hypertension, cardiac arrhythmia)

... and 6 more (see full listing online)

## Locations (4 total)

---

John R. Oishei Children's Hospital of Buffalo, Buffalo, New York, United States

Columbia University Irving Medical Center NewYork-Presbyterian Morgan Stanley Children's Hospital, New York, New York, United States

University of Rochester, Golisano Children's Hospital, Rochester, New York, United States

... and 1 more locations