

Transcutaneous Posterior Tibial Nerve Stimulation for Premature Ejaculation

NCT04207723

Status	RECRUITING
Phase	Not Applicable
Sponsor	Boston Medical Group
Enrollment	120 participants

Key Eligibility Criteria

Inclusion (8)

- Primary premature ejaculation as defined by the International Society for Sexual Medicine (ISSM-International Society for Sexual Medicine) (30): a) ejaculation occurs always or almost always within the first minute after penetration, b) disability to delay ejaculation in all or almost all penetrations, c) negative personal consequences are generated, such as stress, discomfort, frustration and / or avoidance of sexual intimacy.
- Age equal to or greater than 18 years.
- PEDT score greater than 11.
- Stable heterosexual relationship of at least 6 months with the interest of maintaining it for at least the duration of the study.
- Sexual activity at least once a week.
- ... and 3 more (see full listing online)

Exclusion (11)

- IIEF-EF score greater than 25.
- Clinically significant comorbidity: cardiovascular, hepatic, thromboembolic, neurological, locomotive, endocrine, oncological, renal or rheumatologic.
- History of retroperitoneal surgery, radiotherapy or multiple sclerosis.
- History of mental illness: depression, anxiety, suicidal behavior, bipolar disorder, agoraphobia, dysthymia, social phobia, obsessive compulsive disorder, post-traumatic stress, psychiatric disorder, referred by the patient or by the use of a medication for one of these terms.
- Consumption of medications that affect ejaculatory control such as psychiatric medications, opioid analgesics, alpha blockers.
- ... and 6 more (see full listing online)

Locations (1 total)

Boston Medical Group, Bogotá, Bogota D.C., Colombia