

# Ticagrelor With Low-dose Versus Regular Aspirin in Patients With Acute Coronary Syndrome (ACS) at High-Risk for Ischemia After Percutaneous Coronary Intervention

NCT04240834

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 4
<b>Sponsor</b>	Fu Wai Hospital, Beijing, China
<b>Enrollment</b>	1,220 participants

## Key Eligibility Criteria

---

### Inclusion (19)

- ACS patients at high risk for ischemic events after successful PCI with implantation of at least one drug eluting stent
- Able and willing to provide informed consent and participate in 12 months follow-up period
- Able to receive DAPT treatment
- Family history of premature coronary heart disease (Coronary heart disease in first-degree male relative <55 years old or in first-degree female relative <65 years old)
- Repeated myocardial infarction
- ... and 14 more (see full listing online)

### Exclusion (10)

- Need for chronic oral anticoagulation
- With cardiomyopathy HCM/DCM/RCM
- With severe ventricular arrhythmia requiring ICD implantation
- With chronic respiratory disease (COPD, asthma, chronic bronchitis, pulmonary heart disease)
- With severe infectious disease active hepatitis B, active hepatitis C, AIDS
- ... and 5 more (see full listing online)

## Locations (1 total)

---

Fuwai Hospital, Beijing, Beijing Municipality, China