

The Evaluation of Banana Flower Stamens Extract on Prevention of Benign Prostatic Hyperplasia in Adults

NCT04266418

Status	RECRUITING
Phase	Not Applicable
Sponsor	TCI Co., Ltd.
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (6)

- Male aged 40-80 years old
- dPSS score ≤ 19
- The subject did not take α -blocker or anticholinergic agents in the last 8 weeks. The subject did not take 5 α -reductase inhibitor or androgen suppression agents in the last 16 weeks (depending on medical history).
- The subject isn't diagnosed with cancer
- The subject is able to read and finish the information on the questionnaire.
- ... and 1 more (see full listing online)

Exclusion (6)

- The subject has a history of epilepsy or convulsions, liver and kidney disease, cancer, endocrine disease, mental illness, alcohol or drug abuse, and other major organic diseases (depending on medical history).
- The lower urinary tract urination symptoms of the subject are not related to prostatic hypertrophy (depending on medical history).
- Residual urine volume > 250 mL (depending on medical history)
- Subjects have had pelvic radiation therapy or pelvic surgery (including prostate or bladder surgery, but those who only have had a prostate slice can participate in the trial).
- Subjects have taken sexual hormone preparations including LHRH agonists, anti-androgens, feminine, or Penta-reductase inhibitors (Proscar and Avodart) 16 weeks prior to the trial.
- ... and 1 more (see full listing online)

Locations (1 total)

National Taiwan University Hospital, Taipei, Taiwan