

Duration of Urinary Catheterization

NCT04359069

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Southern California
Enrollment	176 participants

Key Eligibility Criteria

Inclusion (6)

- Elective procedures involving total mesorectal excision, including low anterior resection and abdominoperineal resection for rectal cancer as well as proctectomy for inflammatory bowel disease.
- All approaches (open, laparoscopic and robotic) will be included, as the approaches not differ in the total mesorectal excision technique.
- Patients who received neoadjuvant chemotherapy and/or radiation treatments will be included.
- Age e 18 years.
- American Society of Anesthesiologists (ASA) class I-III.

... and 1 more (see full listing online)

Exclusion (11)

- Patients undergoing emergent or urgent surgery.
- Patients undergoing total mesorectal excision in combination with other major surgical procedures on the same day should be excluded from this study.
- Patients sustaining genitourinary tract injury during the procedure should be excluded postoperatively.
- History of urinary retention after previous procedure, surgery, or urinary catheter removal.
- History of urinary retention not being actively treated.

... and 6 more (see full listing online)

Locations (2 total)

Keck Hospital of USC, Los Angeles, California, United States
Los Angeles County Hospital (LAC/USC), Los Angeles, California, United States