

Antibiotic Prophylaxis for Bladder Botox

NCT04444440

Status	RECRUITING
Phase	Phase 4
Sponsor	University of Alberta
Enrollment	202 participants

Key Eligibility Criteria

Inclusion (3)

- Patients with idiopathic OAB diagnosed clinically who have failed medical management
- Female
- Age e 18

Exclusion (7)

- Patients with neurogenic OAB (OAB with potential underlying neurologic cause - multiple sclerosis, spinal cord injury, Parkinson's disease, other)
- Contraindication to injection of Botox - hypersensitivity to any botulinum toxin preparation or to any of the components in the formulation, current urinary tract infection, symptomatic urinary retention or PVR \gt 200 mL, unwillingness or inability to initiate CIC post-treatment if required.
- Contraindication to oral Ciprofloxacin - hypersensitivity or allergy to Ciprofloxacin or other fluoroquinolone, concurrently taking Tizanidine or Agomelatine.¹⁴
- Active antibiotic therapy for any indication at the time of Botox injection - increased risk of adverse reaction with combining antibiotics, reduced risk of UTI with additional antibiotic.
- Male

... and 2 more (see full listing online)

Locations (1 total)

Urogynecology Clinic - Royal Alexandra Hospital, Edmonton, Alberta, Canada