

Combination of Ranibizumab and Targeted Laser Photocoagulation

NCT04444492

Status	RECRUITING
Phase	Phase 3
Sponsor	University of Giessen
Enrollment	110 participants

Key Eligibility Criteria

Inclusion (7)

- Diagnosis of macular edema due to central retinal vein occlusion foveal thickness $\geq 250 \mu\text{m}$ (measured by OCT)
 - Age ≥ 18 years
 - Written informed consent of the patient
 - BCVA score in the study eye between 24 letters (20/320) and 78 letters (20/25) measured in ETDRS chart
 - History of CRVO no longer than 6 months
- ... and 2 more (see full listing online)

Exclusion (18)

- CRVO with ischemic maculopathy defined as diameter of the foveolar avascular zone larger than 2 optic disc diameters
 - Macular edema due to another etiology than retinal vein occlusion (e.g. diabetic maculopathy, uveitis, age related macular degeneration, Irvine-Gass syndrome)
 - History of idiopathic central serous chorioretinopathy
 - Presence of vitreoretinal interface disease (e.g. vitreomacular traction, epiretinal membrane), either on clinical examination or in OCT
 - An eye that, in the investigator's opinion, would not benefit from resolution of macular edema, such as eyes with foveal atrophy, dense pigmentary changes, or dense subfoveal hard exudates
- ... and 13 more (see full listing online)

Locations (15 total)

Medizinische Universität Innsbruck, Klinik für Augenheilkunde und Optometrie, Innsbruck, Austria
Universitätsklinikum Carl Gustav Carus Dresden Klinik und Poliklinik für Augenheilkunde, Dresden, Germany
Internationale Innovative Ophthalmochirurgie GbR, Klinik für Augen Chirurgie, Düsseldorf, Germany
... and 12 more locations