

Diagnostic Role of the "White Test" With Lipidic Solution in the Early Intraoperative Identification of Open Bile Ducts for the Prevention of Bile Leakage After Liver Resection (BiLe -Trial)

NCT04523701

Status	RECRUITING
Phase	Phase 3
Sponsor	Giulia Manzini
Enrollment	210 participants

Key Eligibility Criteria

Inclusion (4)

- Patients who will receive an anatomical resection of two or more liver segments for any reason with simultaneous cholecystectomy in elective Setting
- Patients who will receive an anatomical resection of two or more liver segments for any reason which already had a cholecystectomy if intraoperatively the cystic stump can be identified and opened
- Ability of subject to understand character and individual consequences of the clinical Trial
- Informed consent documented by signature

Exclusion (6)

- Previous cholecystectomy if intraoperative it is not possible to identify the cystic stump
 - Intraoperative hepatico-jejunostomy
 - Hypersensitivity to fish-, egg-, soybean or peanut protein or to any of the active ingredients or excipients
 - Immunosuppression, pregnancy
 - Emergency liver resection because of traumatic liver rupture
- ... and 1 more (see full listing online)

Locations (5 total)

Department of Visceral Surgery, Cantonal Hospital of Aarau, Aarau, Switzerland
Clarunis Basel, Universitäres Bauchzentrum Basel, Basel, Switzerland
Kantonsspital Luzern, Lucerne, Switzerland
... and 2 more locations