

The Effect of Arthroscopic Subacromial Decompression in Patients Who Are Non-responders to Non-operative Treatment.

NCT04644042

Status	RECRUITING
Phase	Not Applicable
Sponsor	Hvidovre University Hospital
Enrollment	160 participants

Key Eligibility Criteria

Inclusion (8)

- SIS diagnosis (Consultant's clinical diagnosis of SIS + at least 3 out of 5 positive tests from the following: Hawkin's, Neer's, Jobe's, Painful arc and external rotation resistance test)
- Positive subacromial injection test
- Insidious onset of shoulder pain
- Considered a surgical candidate by an orthopedic shoulder specialist.
- Symptoms for at least 6 months
- ... and 3 more (see full listing online)

Exclusion (5)

- Terminal illness or severe medical illness (ASA score higher than or equal to 4), systemic musculoskeletal disease, inflammatory joint disease (e.g. rheumatoid arthritis), symptomatic cervical spine pathology or thoracic outlet syndrome.
- Full-thickness rotator cuff tear, calcified tendonitis, labral tear, frozen shoulder, biceps tendon pathology, acromioclavicular osteoarthritis, glenohumeral osteoarthritis or other concomitant shoulder pathology.
- Previous surgery or radiotherapy on the affected shoulder.
- Pregnancy
- Ongoing workers compensation case or job rehabilitation process

Locations (1 total)

Hvidovre Univeristy Hospital, Hvidovre, Denmark, Denmark

<https://clinicaltrials.gov/study/NCT04644042>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).