

Pyloric Sphincter Abnormalities in Patients With Gastroparesis Symptoms

NCT04661215

Status RECRUITING
Sponsor Johns Hopkins Bloomberg School of Public Health
Enrollment 150 participants

Key Eligibility Criteria

Inclusion (14)

- Provision of signed and dated informed consent form.
 - Stated willingness to comply with all study procedures and availability for the duration of the study.
 - Male or female, aged 18-85
 - Symptoms of gastroparesis, either diabetic or idiopathic etiology
 - Symptoms of gastroparesis with minimum Gastroparesis Cardinal Symptom Index (GCSI) score of 2.0
- ... and 9 more (see full listing online)

Exclusion (20)

- Prior gut lumen surgery on the esophagus or the stomach, including Nissen fundoplication.
 - Prior surgery on the pylorus (G-POEM, surgical pyloroplasty, surgical pyloromyotomy)
 - Known history of achalasia or esophageal stricture
 - Known history of physiological or mechanical GI obstruction
 - Abnormalities seen on a prior upper endoscopy placing patient at increased risk:
- ... and 15 more (see full listing online)

Locations (5 total)

Mayo Clinic Arizona, Scottsdale, Arizona, United States
University of Louisville, Louisville, Kentucky, United States
Massachusetts General, Boston, Massachusetts, United States
... and 2 more locations