

# Reduction in Number of Botox Injections for Urgency Urinary Incontinence

NCT04731961

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Virginia
Enrollment	56 participants

## Key Eligibility Criteria

---

### Inclusion (9)

- Females at least 21 years of age
  - English speaking
  - Five or more urge urinary incontinence episodes on a three-day voiding diary. Urge incontinence episodes will be determined based on voiding diary and subject indication of coincident urge symptoms, allowing for self-characterization of incontinence type
  - Urge predominant (urge >50% of total incontinent episodes) urinary incontinence based on self-reported characterization of incontinent episodes on diary
  - Patient or caregiver willing to perform clean intermittent catheterization, or patient willing to have indwelling Foley catheter in the event this would be required
- ... and 4 more (see full listing online)

### Exclusion (20)

- Current symptomatic urinary tract infection that has not resolved prior to randomization.
  - Baseline need for intermittent self-catheterization
  - PVR (Post void residual) >150 mL on 2 occasions with void(s) of greater than 150 mL
  - Surgical treatment for stress incontinence (sling, Burch or urethral injection) or pelvic organ prolapse recommended or planned at enrollment by study investigator(s).
  - Any prior intra-detrusor botulinum toxin A injections
- ... and 15 more (see full listing online)

## Locations (1 total)

---

University of Virginia, Charlottesville, Virginia, United States