

Efficacy of a Right-sided Ablation of the Anterior Ganglionated Plexus for Neurally Mediated Syncope

NCT04755101

Status	RECRUITING
Phase	Not Applicable
Sponsor	Imelda Hospital, Bonheiden
Enrollment	110 participants

Key Eligibility Criteria

Inclusion (3)

- Patients must be in sinus rhythm and have ≥ 3 syncopes during the last 18 months^{*} and a previous positive tilt table test (TTT) with a cardioinhibitory or mixed response (VASIS I, IIA or IIB classification).
- ^{*} syncopes occurring during TTT are not taken into account
- Patients have a 'preserved cholinergic SN reserve', defined as $\geq 20\%$ sinus heart rate increment during a pharmacological test with atropine.

Exclusion (16)

- < 14 years age
- Any unstable medical condition, life expectancy < 12 months
- Inability to provide consent or undergo follow-up
- Syncope due to a non-cardiac disease or due to an advanced neuropathy
- Moderate to severe valvular or subvalvular aortic stenosis or mitral stenosis
- ... and 11 more (see full listing online)

Locations (3 total)

Imeldaziekenhuis, Bonheiden, Belgium
Algemeen Ziekenhuis Sint Jan, Bruges, Belgium
Universitair Ziekenhuis Gasthuisberg, Leuven, Belgium