

Prospective Evaluation of Topical Almond Oil vs Hydroquinone

NCT04875715

Status	RECRUITING
Phase	Phase 4
Sponsor	University of California, Davis
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (1)

- Premenopausal women of Fitzpatrick skin types 3 to 6

Exclusion (13)

- Those with a nut allergy
- Smoking is an independent risk factor and serves as a confounder for the development of facial aging [18]. Therefore current smokers, those that have smoked within the past year, and former smokers with greater than a 20 pack year history of smoking will be excluded.
- Those with an autoimmune photosensitive condition or a known genetic condition with a deficiency in collagen production (such as Ehlers-Danlos syndrome) will be excluded as this can be a confounder for facial photoaging.
- Hormonal medications or therapies will not be allowed unless they have been stable on their medications for at least 2 months. Examples include oral contraceptives, progesterone-based injections, and estrogen based hormonal replacement therapy.
- Those who have undergone any cosmetic procedures to the face in the 4 months prior to enrollment in the opinion of the investigator. People that have undergone deeper chemical peels (TCA based peels) within 1 year prior to enrollment will also not be eligible.

... and 8 more (see full listing online)

Locations (1 total)

UC Davis Department of Dermatology, Clinical Trials Unit, Sacramento, California, United States