

Catheter-Related Early Thromboprophylaxis With Enoxaparin Studies

NCT04924322

Status	RECRUITING
Phase	Phase 2, Phase 3
Sponsor	Yale University
Enrollment	258 participants

Key Eligibility Criteria

Inclusion (3)

- >36 weeks corrected gestational to <17 years old
- <24 hours after insertion of an untunneled CVC
- CVC inserted in the internal jugular or femoral vein

Exclusion (13)

- Radiologic diagnosis of CADVT in the site of insertion in prior 6 weeks
- Currently receiving an antithrombotic agent, e.g., LMWH, UFH, warfarin and aspirin, but not UFH at dose to maintain patency of a vascular catheter
- Presence of clinically relevant bleeding, i.e., hemoglobin decreased e2 g/dl in 24 hours, required medical or surgical intervention to restore hemostasis, or in the retroperitoneum, pulmonary, intracranial or central nervous system, in the prior 60 days
- Surgery in the prior 7 days
- Major trauma in the prior 7 days
- ... and 8 more (see full listing online)

Locations (22 total)

Children's of Alabama, Birmingham, Alabama, United States
Arkansas Children's Hospital, Little Rock, Arkansas, United States
Children's Hospital Colorado, Aurora, Colorado, United States
... and 19 more locations