

Gut Kidney Axis in Enteric Hyperoxaluria

NCT05124886

Status	RECRUITING
Phase	Not Applicable
Sponsor	NYU Langone Health
Enrollment	80 participants

Key Eligibility Criteria

Inclusion (8)

- In order to be eligible to participate in this study, an individual must meet all of the following criteria:
- Subjects \geq 18 years and \leq 80 years of age
- a confirmed diagnosis of IBD or \geq 6 months post- RYGB, with a diagnosis USD event (renal colic with spontaneous stone passage, emergency room visits, or urological interventions) or have kidney stone on imaging (CT, MRI, or US) in the previous three years.
- We will include all racial and ethnic groups, and both men and women.
- In order to be eligible in this study as a healthy control, an individual must meet all of the following criteria:
... and 3 more (see full listing online)

Exclusion (14)

- An individual who meets any of the following criteria will be excluded from participation in this study:
- pregnant or nursing women and people with any medical, psychiatric, debilitating disease/disorder or social condition that, in the judgment of the investigator, would interfere with or serve as a contraindication to adherence to the study protocol, ability to give ICF or complete the protocol.
- subjects with total and partial colectomy.
- subjects who received oral, intramuscular, or intravenous antibiotics within three months before screening.
- Patients with perianal disease usually receive recurrent antibiotics and, therefore, will be excluded from the study.
... and 9 more (see full listing online)

Locations (1 total)

NYU Langone Health, New York, New York, United States