

Study of Gastric Motility in Eosinophilic Gastritis

NCT05229432

Status	RECRUITING
Sponsor	Children's Hospital Medical Center, Cincinnati
Enrollment	30 participants

Key Eligibility Criteria

Inclusion (4)

- Eosinophilic gastritis (EG) defined as at least one endoscopy with histopathologic evidence of ≥ 30 eosinophils in 5 or more high powered fields (hpf's) with associated symptoms of EG.
- Patient reported symptoms starting at least one year or more prior to screening consistent with a diagnosis of EG: nausea, vomiting, early satiety, abdominal pain/bloating, regurgitation, diarrhea.
- Symptoms suggestive of a possible gastric motility disorder during the 60 days prior to screening including: nausea, vomiting, early satiety, post-prandial feeling of fullness or bloating.
- Tolerance and willingness to consume the oatmeal meal in this study.

Exclusion (8)

- Diagnosis of eosinophilic enteritis.
- Inflammatory bowel disease (e.g., Crohn's disease or ulcerative colitis).
- Known disorders associated with eosinophilic gastrointestinal (GI) diseases (e.g., D 816 V Kit+ systemic mastocytosis, Marfan syndrome or Loey's Dietz Syndrome).
- Known inflammatory or autoimmune disorders associated with gastric dysmotility such as systemic sclerosis, lupus or eosinophilic fasciitis.
- Uncontrolled diabetes or known diabetic complications of gastroparesis, neuropathy or nephropathy.

... and 3 more (see full listing online)

Locations (3 total)

Northwestern University, Chicago, Illinois, United States
The National Institutes of Health, Bethesda, Maryland, United States
University of Utah, Salt Lake City, Utah, United States