

# Ketorolac in Upper Extremity Tendinopathy and Arthropathy

NCT05292339

---

Status	RECRUITING
Phase	Phase 4
Sponsor	Emory University
Enrollment	160 participants

## Key Eligibility Criteria

---

### Inclusion (3)

- Age 18 or older,
- symptomatic tendinopathy or arthropathy of the shoulder, elbow, hand, or wrist who have not undergone prior surgical treatment for their condition.
- Diagnoses may include trigger finger, De Quervain's tenosynovitis, radiocarpal osteoarthritis, first carpometacarpal (CMC) joint osteoarthritis, metacarpophalangeal joint osteoarthritis, or proximal interphalangeal joint osteoarthritis.

### Exclusion (5)

- Patients under the age of 18,
- Patients who have undergone prior triamcinolone or ketorolac injections within the past 6 months,
- Patients who have undergone prior surgical treatment for their hand condition,
- Patients with allergy or contraindication to triamcinolone or ketorolac injection,
- Patients with an active infection at the treatment site \[active infection defined as cellulitis, purulence, fever, chills, or presence of elevated inflammatory markers, ie. white blood cells (WBC), erythrocyte sedimentation rate (ESR), C-reactive protein (CRP)\].

## Locations (4 total)

---

Emory Saint Joseph's Hospital, Atlanta, Georgia, United States  
Emory University Orthopaedic and Spine Center, Atlanta, Georgia, United States  
12 Executive Park Drive, Atlanta, Georgia, United States  
... and 1 more locations