

Trazodone vs. Quetiapine for the Treatment of ICU Delirium

NCT05307003

Status	RECRUITING
Sponsor	University of Southern California
Enrollment	60 participants

Key Eligibility Criteria

Inclusion (4)

- ≥ 18 -years-old
- Admitted to the surgical ICU for > 24 hours
- Diagnosis of ICU delirium defined by positive CAM-ICU score AND exhibiting symptomatic delirium (i.e., combative, pulling at lines, a danger to self or others, inability to sleep, hallucinations, etc.), thus, requiring the need for pharmacologic intervention as determined by the attending intensivist
- Receiving either quetiapine or trazodone for the treatment of delirium

Exclusion (28)

- Presence of an acute neurologic condition (i.e., acute cerebrovascular accident, intracranial tumor, traumatic brain injury, etc.) on ICU admission. History of stroke or other neurological condition(s) without cognitive impairment is not an exclusion criterion.
- Pregnancy/lactation
- History of ventricular arrhythmia including torsade de pointes or second- and third-degree heart block
- Allergy/hypersensitivity reaction to trazodone and/or quetiapine
- Diagnosis of dementia
- ... and 23 more (see full listing online)

Locations (1 total)

Keck Hospital of the University of Southern California, Los Angeles, California, United States