

To Evaluate the Therapeutic Effect of Laser Acupuncture in Patients With Acute or Subacute Post Stroke Shoulder Pain

NCT05328401

Status	RECRUITING
Phase	Not Applicable
Sponsor	Chang Gung Memorial Hospital
Enrollment	58 participants

Key Eligibility Criteria

Inclusion (6)

- Beyond the age of 20.
- Normal consciousness before and after stroke who can reply and fill in the questionnaire.
- Subacute (subacute phase: 7 days to 6 months after stroke) and Chronic (chronic phase: more than 6 months after stroke) with shoulder pain after stroke and in post- stroke recovery period.
- Stop taking painkillers for more than 1 day,
- Without received intramuscular drugs (steroids, botulinum toxin, glucose, etc.) injections in the shoulder joint or around the shoulder for more than 2 weeks
- ... and 1 more (see full listing online)

Exclusion (9)

- Participants with shoulder pain symptoms before the stroke
- Acute shoulder trauma on the affected side of shoulder pain in the past.
- Surgery on the affected side of shoulder pain in the past.
- Those who cannot complete the scale assessment due to aphasia, post-stroke depression, and obvious cognitive impairment.
- Women who are pregnant or planning to become pregnant.
- ... and 4 more (see full listing online)

Locations (1 total)

Chang Gung Memorial Hospital, Keelung Medical Center, Keelung, Taiwan, Taiwan