

Does Additional Use of Preoperative Azithromycin Decrease Posthysterectomy Infections

NCT05337566

Status	RECRUITING
Phase	Not Applicable
Sponsor	Helsinki University Central Hospital
Enrollment	2,278 participants

Key Eligibility Criteria

Inclusion (1)

- Women undergoing hysterectomy for benign indication in University Hospitals (Helsinki University Hospital, Turku University Hospital, Tampere University Hospital, Oulu University Hospital and Kuopio University Hospital) who have not any contraindications for azithromycin or cefuroxime.

Exclusion (6)

- Inability to understand the study protocol.
- Allergy for either cefuroxime or azithromycin.
- Congenital or acquired prolonged Q-T-corrected interval. All the participants will be asked about arrhythmias and whether they have congenital arrhythmias in the family,
- Electrocardiogram will be checked for all the participants.
- Use of medicines that may prolong Q-T-corrected interval (class Ia arrhythmia medications, quinidine, procainamide, and class III arrhythmia medications dofetilide, amiodarone and sotalol).

... and 1 more (see full listing online)

Locations (6 total)

Helsinki University Central Hospital, Helsinki, Finland

Jyväskylä Central Hospital, Jyväskylä, Finland

Kuopio University Central Hospital, Kuopio, Finland

... and 3 more locations