

FotonaSmooth Erbium:YAG Laser for the Treatment of Urethral Pain Syndrome in Women

NCT05344716

Status	RECRUITING
Phase	Not Applicable
Sponsor	Prof. Dr. Volker Viereck
Enrollment	30 participants

Key Eligibility Criteria

Inclusion (5)

- Adult female, 18 years of age or older
- Urethral pressure pain upon palpation, VAS Score e 2
- Intermittent or chronic pain in the urethra and/or the small pelvis (independent of micturition) for at least 6 months
- No significant improvement of UPS from at least one previous conservative treatment (pessary, antibiotics, local oestrogens etc.)
- Signed informed consent

Exclusion (12)

- Pregnancy
- Treatment with Isotretinoin (Acne, Rosacea) within last 6 months
- Acute urinary tract infection or other acute infection of the bladder, vagina, vulva or urinary tract detected by routine urine analysis
- Positive urethral swab for Ureaplasma, Mycoplasma or Chlamydia d 6 weeks. Exception: Patients with chronic and persistent fastidious bacteria can be included six weeks after two unsuccessful treatments with antibiotics. Antibiotics therapy must also include sexual partners.
- Pre-existing bladder or urethra pathology

... and 7 more (see full listing online)

Locations (1 total)

Blasenzentrum der Frau AG, Frauenfeld, Thurgau, Switzerland