

Prevention of Postpartum Hemorrhage With Tranexamic Acid

NCT05370820

Status	RECRUITING
Phase	Phase 2
Sponsor	Inova Health Care Services
Enrollment	120 participants

Key Eligibility Criteria

Inclusion (5)

- Women who are scheduled to under medically indicated cesarean section at greater than 34+0 weeks gestation or women who are scheduled to undergo an elective cesarean section at 39+0 weeks gestation in accordance with recommendations from the American Congress of Obstetricians and Gynecologists
- Women who are indicated to have a vaginal delivery at $> 34+0$ weeks gestation.
- Pregnant women with normal serum creatinine (serum creatinine < 0.9) within 2 weeks of estimated/scheduled delivery
- Women between the ages of 18 and 50 years old
- Ability to understand and the willingness to sign a written informed consent form and HIPAA Authorization.

Exclusion (10)

- active thrombotic or thromboembolic disease
- a history of arterial or venous thromboembolic event
- inherited thrombophilia or preexisting conditions that predisposes them to thromboembolic events (i.e. lupus, antiphospholipid syndrome, thrombocytosis or thrombophilic thrombocytopathy)
- a subarachnoid hemorrhage
- acquired defective color vision
- ... and 5 more (see full listing online)

Locations (2 total)

George Washington University Hospital, Washington D.C., District of Columbia, United States
Inova Fairfax Medical Campus, Falls Church, Virginia, United States