

Efficacy of Surgical Joint Denervation in Painful Digital Osteoarthritis

NCT05378841

Status	RECRUITING
Phase	Not Applicable
Sponsor	Assistance Publique - Hôpitaux de Paris
Enrollment	19 participants

Key Eligibility Criteria

Inclusion (7)

- Age e 18 years
 - Patients with symptomatic digital osteoarthritis according to the criteria of the American College of Rheumatology
 - painful PIP joint (Analog digital scale e 4/10) for more than 3 months
 - having failed the usual medical treatment for a minimum of 3 months (level 1 analgesics, NSAIDs, infiltrations)
 - Signature of informed consent
- ... and 2 more (see full listing online)

Exclusion (10)

- digital osteoarthritis secondary to other known causes (eg gout, reactive arthritis, RA, psoriatic arthritis, spondyloarthropathies, septic arthritis)
 - anesthetic contraindications to surgery or to performing an MRI
 - Existence of a pain syndrome in the upper limbs, which would interfere with the assessment of AD.
 - Fibromyalgia
 - Use of oral, intramuscular or intra-articular or intravenous corticosteroids, immunosuppressants (methotrexate, sulfasalazine, leflunomide, biomedicines), hyaluronic acid infiltration in the finger joints in the previous month
- ... and 5 more (see full listing online)

Locations (1 total)

Rheumatology department Saint-Antoine Hospital, Paris, France