

Washed Microbiota Transplantation for Malnutrition After Nonphysiological Reconstruction of the Upper Gastrointestinal Tract

NCT05439135

Status	RECRUITING
Phase	Not Applicable
Sponsor	The Second Hospital of Nanjing Medical University
Enrollment	62 participants

Key Eligibility Criteria

Inclusion (4)

- Patients undergone nonphysiological reconstruction of the gastrointestinal tract (such as Billroth-II, Roux-en-Y, Child, Whipple reconstruction)
- Aged over 18 years old;
- Malnutrition is assessed by Nutritional Risk Screening 2002 score³ and Patient Generated Subjective Global Assessment (PG-SGA) score⁴;
- Written informed consent;

Exclusion (14)

- Expected survival time < 3 months;
- Still on anti-tumor therapy or it is expected during the visit;
- Complicated with ascites or edema caused by malnutrition;
- Known organic gastrointestinal disease (e.g., gastrointestinal infection, inflammatory bowel disease, or radiation enteritis);
- Undergone other gastrointestinal surgery (e.g., enterectomy, enterostomy, or gastrostomy) except appendectomy or anal fistula surgery;
- ... and 9 more (see full listing online)

Locations (1 total)

The Second Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu, China