

Intracoronary Stenting and Restenosis - Randomized Trial of Drug-eluting Stent Implantation or Drug-coated Balloon Angioplasty According to Neointima Morphology in Drug-eluting Stent Restenosis 5

NCT05544864

Status	RECRUITING
Phase	Not Applicable
Sponsor	Deutsches Herzzentrum Muenchen
Enrollment	376 participants

Key Eligibility Criteria

Inclusion (5)

- Patients with ischemic symptoms and/or evidence of myocardial ischemia
- Presence of e 50% restenosis after prior implantation of drug-eluting stents in native coronary vessels.
- Availability of an OCT-pullback of the target lesion
- Written informed consent by the patient for participation in the study.
- Age e 18 years

Exclusion (11)

- Cardiogenic shock
- Acute ST-elevation myocardial infarction within 48 hours from symptom onset.
- Target lesion located in left main trunk or bypass graft.
- Additional coronary intervention planned within 30 days of the procedure.
- Non-successful treatment of other lesion(s) during the same procedure
- ... and 6 more (see full listing online)

Locations (5 total)

Universitätsklinikum Frankfurt, Frankfurt am Main, Hesse, Germany
Elisabeth-Krankenhaus Essen GmbH, Essen, Germany
Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany
... and 2 more locations