

# Durg Coated Balloon Angioplasty in Infrapopliteal Lesions

NCT05620095

---

<b>Status</b>	RECRUITING
<b>Sponsor</b>	Liyuan Hospital of Tongji Medical College, Huazhong University of Science and Technology
<b>Enrollment</b>	1,000 participants

## Key Eligibility Criteria

---

### Inclusion (9)

- Rutherford grade 4-6.
- Patients who understand the purpose of this study, volunteer to participate in the experiment, sign informed consent and are willing to follow up.
- Single or sequential de novo or restenotic lesions (stenosis e 70% diameter reduction or occlusion) in the infrapopliteal arteries \>20 mm. Lesions should not extend beyond the ankle joint.
- Successful wire crossing of the lesion. After the pre-dilation of the ordinary balloon, the angiography showed that there was continuous blood flow.
- At least one of the infrapopliteal arteries received a drug-coated balloon.

... and 4 more (see full listing online)

### Exclusion (7)

- Blood flow was not successfully reestablished.
- Patients with stroke, cerebral hemorrhage, gastrointestinal hemorrhage or myocardial infarction within 3 months before enrollment.
- Patients who are known to be allergic to heparin, aspirin, other antiplatelet drugs, contrast agents, etc.
- Patients who have participated in clinical trials of drugs or other medical devices that interfere with this clinical trial within the past 3 months.
- Pregnant and lactating women.

... and 2 more (see full listing online)

## Locations (10 total)

---

Xuanwu Hospital Capital Medical University, Beijing, Beijing Municipality, China

Zibo Feng, Wuhan, Hubei, China

the second Affiliated Hospital of Medical College of Suzhou University, Suzhou, Jiangsu, China

... and 7 more locations