

# Best Antithrombotic Therapy in Patients With Acute Venous Thromboembolism While Taking Antiplatelets

NCT05627375

---

Status	RECRUITING
Phase	Phase 3
Sponsor	Centre Hospitalier Universitaire de Saint Etienne
Enrollment	1,400 participants

## Key Eligibility Criteria

---

### Inclusion (6)

- Signed informed consent
  - Patients with acute objectively confirmed symptomatic proximal deep-vein thrombosis (DVT) or pulmonary embolism (PE) (with or without deep-vein thrombosis). Proximal deep-vein thrombosis is defined as thrombosis involving at least the popliteal vein or a more proximal vein of the lower limb.
  - Indication of full-dose anticoagulant therapy for at least 3 months.
  - Prescription of antiplatelet therapy for secondary prevention of atherosclerotic cardiovascular diseases, at the time of VTE diagnosis
  - Life expectancy more than 3 months
- ... and 1 more (see full listing online)

### Exclusion (9)

- Unable to give informed consent
  - Active bleeding or a high risk of bleeding contraindicating anticoagulant treatment; a systolic blood pressure of more than 180 mm Hg or a diastolic blood pressure of more than 110 mm Hg
  - Anticoagulation for more than 5 days prior to randomization
  - Active pregnancy or expected pregnancy or no effective contraception
  - Isolated distal deep vein thrombosis
- ... and 4 more (see full listing online)

## Locations (28 total)

---

CHU Amiens, Amiens, France  
CHU Angers, Angers, France  
CHU Besançon - Hôpital Jean Minjot, Besançon, France  
... and 25 more locations

---

<https://clinicaltrials.gov/study/NCT05627375>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).