

A Phase 2 Trial Comparing Antiviral Treatments in Early Symptomatic Influenza

NCT05648448

Status	RECRUITING
Phase	Phase 2
Sponsor	University of Oxford
Enrollment	3,000 participants

Key Eligibility Criteria

Inclusion (6)

- Patient understands the procedures and requirements and is willing and able to give informed consent for full participation in the study
 - Adults, male or female, aged 18 to 60 years at time of consent.
 - Early symptomatic Influenza (A or B); at least one reported symptom of influenza (including fever, history of fever, myalgias, headache, cough, fatigue, nasal congestion, rhinorrhoea and sore throat) within 4 days (96 hours)
 - Influenza positive by rapid antigen test OR a positive RT-PCR test for influenza viruses within the last 24hrs with a Ct value of ≤ 30
 - Able to walk unaided and unimpeded in activities of daily living (ADLs)
- ... and 1 more (see full listing online)

Exclusion (16)

- The patient may not enter the study if ANY of the following apply:
 - Taking any concomitant medications or drugs which could interact with the study medications or have antiviral activity
 - Presence of any chronic illness/condition requiring long term treatment or other significant comorbidity
 - BMI ≥ 35 Kg/m²
 - Clinically relevant laboratory abnormalities discovered at screening
- ... and 11 more (see full listing online)

Locations (4 total)

Universidade Federal de Minas Gerais, Minas Gerais, Brazil
Laos-Oxford-Mahosot Wellcome Trust Research unit, Vientiane, Laos
Sukraraj Tropical & Infectious Disease Hospital, Kathmandu, Nepal
... and 1 more locations

<https://clinicaltrials.gov/study/NCT05648448>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).