

Evaluation of the Benefit of Lidocaine on the Prevention of the Risk of Post Endoscopic Retrograde Cholangio-pancreatography Pancreatitis.

NCT05667987

Status	RECRUITING
Phase	Not Applicable
Sponsor	GCS Ramsay Santé pour l'Enseignement et la Recherche
Enrollment	1,800 participants

Key Eligibility Criteria

Inclusion (3)

- Patient over 18 years of age
- Patient who has read and signed the consent form for participation in the study
- Patient candidate for ERCP with virgin papilla

Exclusion (8)

- Patient with sphincterotomized papilla
 - Patient under court protection, guardianship or curatorship
 - Pregnant woman or woman of childbearing age without highly effective contraception for the duration of the study (surgically sterile, intrauterine device (> 14 days), hormonal contraception (same dose and formulation for at least 6 months), sexual abstinence. Women after menarche and until they become postmenopausal, unless they are permanently infertile or have undergone surgical sterilization, are considered to be of childbearing age, i.e. fertile. A postmenopausal condition is defined as the absence of menstruation for 12 months without any other medical cause
 - Patient not affiliated with the French social security system
 - Patient participating in another clinical research protocol
- ... and 3 more (see full listing online)

Locations (1 total)

Hôpital Privé des Peupliers, Paris, France