

Icodextrin Postpones the Shift of Low Dose to Full Dose Dialysis in the First Year of Incremental Peritoneal Dialysis

NCT05721404

Status	RECRUITING
Phase	Phase 4
Sponsor	First Affiliated Hospital, Sun Yat-Sen University
Enrollment	194 participants

Key Eligibility Criteria

Inclusion (9)

- Age \geq 18 years old;
 - The estimated glomerular filtration rate (eGFR) \geq 3ml/min \cdot 1.73m² when enrolled;
 - The 24-hour urine volume \geq 500ml when enrolled;
 - The patient or his lawful representative is able to receive and complete training of home peritoneal dialysis;
 - Patients were able to follow the follow-up schedule and other requirements of the study;
- ... and 4 more (see full listing online)

Exclusion (13)

- Treated with both peritoneal dialysis and hemodialysis;
 - Patients who have previously received a kidney transplant and have been receiving immunosuppressive therapy;
 - Contraindications to bioelectrical impedance analysis (BIA) testing (e.g., amputation, use of a pacemaker or prosthesis);
 - Allergy to Icodextrin, starch and starch products, or suffer from glycogen storage disease;
 - Contraindications for the use of icodextrin;
- ... and 8 more (see full listing online)

Locations (1 total)

The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China