

SCI-Pex Study - Safety and Performance of PexyEazy®, a Device for Treatment of Hemorrhoids

NCT05782010

Status	RECRUITING
Phase	Not Applicable
Sponsor	Developeration AB
Enrollment	35 participants

Key Eligibility Criteria

Inclusion (3)

- Adult
- Hemorrhoidal disease grade II and III
- American Society of Anesthesiologists (ASA) classification I-III.

Exclusion (15)

- Previous surgical treatment for hemorrhoids by all methods except rubber band ligation or sclerosing injection therapy within the last 3 years.
- Previous surgical treatment with rubber band ligation or injection therapy within the last 2 months.
- More than three surgical treatments with rubber band ligation or sclerosing injection therapy within the last 3 years.
- Under medication with immunosuppressive drugs and/or anticoagulant drugs (not including 75mg acetylsalicylic acid) and not able to interrupt the anticoagulant drug medication or replace it with low molecular weight heparin for 5 days.
- Have ulcerative colitis, Crohn's disease or other intestinal inflammatory disease or presence of inflamed or thickened mucosa of the rectum.

... and 10 more (see full listing online)

Locations (3 total)

Frölunda Specialistsjukhus, Frölunda, Sweden
Norrtälje Sjukhus, Norrtälje, Sweden
Capio kirurgkliniken, Sophiahemmet, Stockholm, Sweden