

Intravitreal Faricimab Injections or Fluocinolone Acetonide (0.19 mg) Intravitreal Implants vs Observation for Prevention of VA Loss Due to Radiation Retinopathy

NCT05844982

Status	RECRUITING
Phase	Phase 3
Sponsor	Jaeb Center for Health Research
Enrollment	600 participants

Key Eligibility Criteria

Inclusion (6)

- Primary uveal melanoma (excluding iris melanoma) receiving primary treatment with plaque brachytherapy
 - Absence of unrelated cause of visual loss
 - Baseline visual acuity \geq 34 letters (20/200 Snellen equivalent or better)
 - Posterior tumor margin \geq 0 mm from the center of the macula (i.e., tumor is NOT under the geometric center of the fovea)
 - Posterior tumor margin \geq 0 mm from the closest disc margin (i.e., tumor is not touching the edge of the optic disc)
- ... and 1 more (see full listing online)

Exclusion (6)

- Opaque media
 - Inability to undergo fluorescein angiography
 - Less than 18 years of age
 - Prior vitrectomy
 - Intraocular pressure (IOP) \geq 25 mmHg or history of steroid- induced IOP elevation that required treatment at baseline
- ... and 1 more (see full listing online)

Locations (22 total)

Byers Eye Institute at Stanford University, Palo Alto, California, United States
Retina Consultants, Manchester, Connecticut, United States
Retina Associates of Florida, LLC, Tampa, Florida, United States
... and 19 more locations