

Study on Prognosis of Acutely Ruptured Intracranial Aneurysms

NCT05851989

Status RECRUITING
Sponsor Haaglanden Medical Centre
Enrollment 880 participants

Key Eligibility Criteria

Inclusion (4)

- Confirmed diagnosis of subarachnoid haemorrhage on CT-scan or lumbar puncture (in the presence of a negative CT-scan)
- Intracranial aneurysm proven within 6 months to be the cause of subarachnoid haemorrhage
- Age 18 years or over at presentation.
- Written informed consent

Exclusion (5)

- Subarachnoid haemorrhage deemed most likely of 'perimesencephalic' origin after consideration of history, clinical examination and radiological findings (including angiographic imaging)
- Subarachnoid haemorrhage deemed most likely of post-traumatic origin after consideration of history, clinical examination and radiological findings (including angiographic imaging)
- Diagnosis of intracerebral arteriovenous malformations or dural arteriovenous fistula.
- No diagnosis of intracranial aneurysm at 6 months after onset of symptoms.
- Not mastering the Dutch language

Locations (6 total)

Amsterdam UMC, Amsterdam, Netherlands
Maastricht UMC, Maastricht, Netherlands
Radboudumc, Nijmegen, Netherlands
... and 3 more locations