

Onlay Synthetic Bioabsorbable Mesh Herniorrhaphy Versus Herniorrhaphy Only in the Primary Treatment of Large Hiatal Hernia

NCT05867225

Status	RECRUITING
Phase	Not Applicable
Sponsor	University Hospital, Bordeaux
Enrollment	256 participants

Key Eligibility Criteria

Inclusion (5)

- Patient undergoing laparoscopic primary repair for symptomatic LHH. LHH is defined as a hiatal hernia larger than 5 cm in axial length as diagnosed at barium series and/or CT-scan with contrast swallow and/or CT-scan with contrast injection with axial reconstruction.
- Symptomatic hiatal hernia: heartburn and/or regurgitation and/or epigastric pain and/or dysphagia and/or anemia (if exclusion of other origin);
- Patient aged \geq 18 years;
- Patient affiliated to a social security system or beneficiary of the same;
- Free written informed consent signed by the participant and the investigator (no later than the inclusion day and before performing any examination required for the study).

Exclusion (11)

- Patient undergoing reoperation for recurrent LHH repair;
 - Emergency presentation needing an operation in a delay $<$ 6 hours;
 - Asymptomatic hiatal hernia;
 - American anesthesiologist score $>$ 3;
 - Recurrent hiatal hernia and previous surgical interventions involving gastroesophageal junction;
- ... and 6 more (see full listing online)

Locations (26 total)

CH de la Côte Basque, Bayonne, France
Hôpital Jean Minjoz, CHRU de Besançon, Besançon, France
Hôpital Cavale blanche, CHRU de Brest, Brest, France
... and 23 more locations