

Impact of Forced Diuresis on the Residual Fragment Rate After Flexible Ureteroscopy for Destruction of Kidney Stones With Laser

NCT05916963

Status	RECRUITING
Phase	Phase 3
Sponsor	University Hospital, Tours
Enrollment	374 participants

Key Eligibility Criteria

Inclusion (5)

- Patients e 18 years old and \< 80 years old
- With the need to perform a flexible ureteroscopy with destruction of the kidney stones with laser
- Participants covered by or entitled to social security
- Written informed consent obtained from the participant
- Ability for participant to comply with the requirements of the study

Exclusion (12)

- Persons covered by articles L1121-5 to L1121-8 of the CSP (corresponding to all protected persons: pregnant women, parturients, nursing mothers, persons deprived of their liberty by judicial or administrative decision, minors, and persons subject to a legal protection measure: guardianship or trusteeship)
- Contra-indication to Furosemide:
- Hyper-sensitivity to the active substance or one of the excipients
- Hyper-sensitivity to Sulfonamide
- Renal failure with oligo-anuria refractory to Furosemide
- ... and 7 more (see full listing online)

Locations (9 total)

CHU d'Angers, Angers, France
Hospices Civils de Lyon, Lyon, France
CHU de Nice, Nice, France
... and 6 more locations