

# Safety and Efficacy of Anakinra Treatment for Patients With Post Acute Covid Syndrome

NCT05926505

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 2, Phase 3
<b>Sponsor</b>	Hellenic Institute for the Study of Sepsis
<b>Enrollment</b>	182 participants

## Key Eligibility Criteria

---

### Inclusion (9)

- Age equal to or above 18 years
  - Male or female gender
  - In the case of women of childbearing age and men, an adequate method of contraception should be used during the study. Contraception should be maintained for at least a period of 3 months after the discontinuation of treatment. As an adequate method of contraception, it is suggested: -male or female condom with or without spermicide -contraceptive cap, a diaphragm or contraceptive sponge with a spermicide Prior to admission to the study, a pregnancy test will be performed to exclude pregnancy to women of childbearing age.
  - Written informed consent provided by the patient. For subjects without decision-making capacity, informed consent must be obtained from a legally designated representative following the national legislation in the Member State where the trial is planned.
  - History of confirmed COVID-19 infection the last 90 days or more
- ... and 4 more (see full listing online)

### Exclusion (16)

- Age below 18 years
  - Denial for written informed consent
  - Any stage IV malignancy
  - Any primary immunodeficiency
  - Less than 1,500 neutrophils/mm<sup>3</sup>
- ... and 11 more (see full listing online)

## Locations (24 total)

---

Out-patient long-COVID department, Jena University Hospital, Jena, Germany  
Out-patient long-COVID department, Patras University General Hospital, Pátrai, Achaia, Greece  
Out-patient long-COVID department III, Evangelismos Athens General Hospital, Athens, Attica, Greece  
... and 21 more locations