

Results of the Use of Two Stentriever Simultaneously Compared With One as a Primary Treatment in Acute Ischemic Stroke

NCT05930145

Status	RECRUITING
Phase	Not Applicable
Sponsor	Fundación EPIC
Enrollment	200 participants

Key Eligibility Criteria

Inclusion (7)

- A new disabling focal neurological deficit compatible with acute cerebral ischemia.
 - Any age. Informed consent obtained from the patient or representative.
 - NIHSS score e 6.
 - Pre-existing functional clinical status less than or equal to 2 according to the mRS clinical scale.
 - Maximum time of 24 hours from symptom onset to arterial puncture. • TICI 0-1 in the diagnosed TICA (terminal internal carotid artery) , MCA (middle cerebral artery), and BA(basilar artery confirmed by angioCT and angiography).
- ... and 2 more (see full listing online)

Exclusion (13)

- Those described in the usual protocols for mechanical thrombectomy of each hospital.
 - Pre-existing functional clinical status greater than 2 according to the mRS clinical scale
 - Patients with tandem lesions of dissection or arteriosclerotic origin located in the extracranial internal carotid artery.
 - Initiation of treatment with a different technique than the one described.
 - Inability to use a proximal balloon guide catheter.
- ... and 8 more (see full listing online)

Locations (4 total)

Alfried Krupp Hospital Ruettenscheid, Essen, Germany
Hospital Universitario de Cruces, Barakaldo, Spain
Hospital Universitario Central de Asturias, Oviedo, Spain
... and 1 more locations