

Early Initiated Vasopressor Therapy in the Emergency Department

NCT05931601

| | |
|------------|----------------------------|
| Status | RECRUITING |
| Phase | Phase 3 |
| Sponsor | Odense University Hospital |
| Enrollment | 320 participants |

Key Eligibility Criteria

Inclusion (1)

- At least 18 years of age

Exclusion (9)

- SBP \leq 100mmHg or MAP \leq 65 mmHg combined with lactate $>$ 2.0 mmol/L,
- Physician defined blood pressure for the individual patient combined with a lactate $>$ 2.0 mmol/L
- Either SBP \leq 100mmHg or MAP \leq 65mmHg with obvious signs of shock with any lactate level evaluated by either two non-specialist physicians (e.g. registrar medical doctors) or a specialist physician.
- Received at least 500ml of intravenous fluid before study inclusion (Including prehospital administration) within the first 4 hours of ED arrival.
- Clinical Frailty Score (CFS) of \leq 4. If CFS is \geq 5 and the treating physician find the patient suitable for ICU admittance, the participant can be enrolled, if the on-call ICU doctor would accept the patient for ICU admittance. If the treating physician is unsure of ICU eligibility, regardless of CFS score, the patient should be consulted with the ICU consultant before study inclusion.

... and 4 more (see full listing online)

Locations (8 total)

Bispebjerg Hospital, Copenhagen, Denmark
Esbjerg Hospital, Esbjerg, Denmark
Gødstrup Regional Hospital, Herning, Denmark
... and 5 more locations

<https://clinicaltrials.gov/study/NCT05931601>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).