

Mycophenolate Mofetil in Gastrointestinal Henoch-Schonlein Purpura

NCT05951517

Status	RECRUITING
Phase	Not Applicable
Sponsor	Sun Yat-Sen Memorial Hospital of Sun Yat-Sen University
Enrollment	24 participants

Key Eligibility Criteria

Inclusion (4)

- Ages range from 3 to 18 years old;
- Patients who meet the diagnostic criteria for HSP and involve the digestive tract: acute diffuse abdominal colic, including intussusception and gastrointestinal bleeding, appears during the course of the disease or is found on physical examination;
- Refractory gastrointestinal type HSP: treated with Intravenous corticosteroid 2 mg/kg/d for 3 days without efficacy or relapsed during the attenuation of corticosteroid
- Parents or guardians agree to treatment and sign a written informed consent form.

Exclusion (5)

- Patients who are allergic to MMF;
- Patients with severe diseases: such as systemic malignancies, heart failure, liver and renal failure, Immune deficiency, severe infectious diseases, organ transplant surgery, or other current indications for emergency surgery;
- Patients with other digestive system diseases;
- Those who have previously used MMF clinical trials;
- Other situations the researcher thought inappropriate to participate in the study.

Locations (1 total)

Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China