

Pituitary Function After Recovery From Septic Shock Among ICU Survivors

NCT05990491

Status RECRUITING
Sponsor Sanjay Gandhi Postgraduate Institute of Medical Sciences
Enrollment 90 participants

Key Eligibility Criteria

Inclusion (7)

- Septic shock group:
- Patients 18-80 years of age who meet the definition of septic shock.
- Vasopressor requirement should be maintained for a period \geq 24 hours and should require ICU stay for a duration of \geq 7 days.
- Patient should recover from shock and be planned for discharge from the ICU
- Non-septic shock group
- ... and 2 more (see full listing online)

Exclusion (7)

- Patients who refuse to provide consent.
- Age $<$ 18 years or $>$ 80 years of age.
- Pregnancy or immediate post-partum ($<$ 6 months post-delivery).
- Chronic kidney disease (Stage 5), chronic liver disease (CHILD B or C), severe Chronic obstructive pulmonary disease, Chronic heart failure.
- Patients with pre-existing hypopituitarism on replacement.
- ... and 2 more (see full listing online)

Locations (2 total)

Department of Critical Care Medicine, SGPGIMS, Lucknow, Uttar Pradesh, India
Department of Endocrinology, SGPGIMS, Lucknow, Uttar Pradesh, India