

# Head Acupuncture Treat Residual Symptoms After Canalith Repositioning Procedure for BPPV

NCT06001047

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Xi'an No.3 Hospital
Enrollment	120 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- According to the diagnostic criteria and therapeutic efficacy evaluation for benign paroxysmal positional vertigo (BPPV) formulated by the Otolaryngology-Head and Neck Surgery Branch of the Chinese Medical Association, the patient was confirmed as having BPPV and the repositioning maneuver was successful.
- After the maneuver, the patient's clinical symptoms were relieved, with no sense of rotation or dizziness, but residual symptoms such as discomfort and unsteady gait persisted.
- There were no obvious communication barriers or visual impairments.
- The patient and their family members agreed to the treatment plan.

### Exclusion (3)

- Patients with other ear diseases, history of head trauma or surgery;
- Patients who are weak and unable to tolerate, pregnant women;
- Patients who have undergone vestibular rehabilitation training before this treatment; c Patients with joint diseases that affect balance and walking; d Patients with a history of brain organic diseases (such as tumors, stroke, cerebral hemorrhage) or severe neurological diseases; e Patients with incomplete data or who withdraw from treatment midway.

## Locations (1 total)

---

Xi'an No3 Hospital, Xi'an, China