

Century Clot-Guided Prophylactic Rivaroxaban for Post STEMI Complicating Left Ventricular Thrombus

NCT06013020

Status	RECRUITING
Phase	Phase 4
Sponsor	Zunyi Medical College
Enrollment	374 participants

Key Eligibility Criteria

Inclusion (2)

- Ischemic chest discomfort for at least 30 minutes, with at least 1-mm (0.1-mv) ST-segment elevation in anterior leads on a standard 12-lead electrocardiogram.
- Patients provide written informed consent prior to enrollment.

Exclusion (9)

- Intracranial, gastrointestinal, or urogenital bleeding within 6 months
- Requiring OAC therapy (eg, atrial fibrillation, deep vein thrombosis, pulmonary thromboembolism);
- Bleeding diathesis, thrombocytopenia (platelet $<100,000/\text{mL}$) or hemoglobin $<10 \text{ g/dL}$, and CRUSADE score-based high bleeding risk
- Hepatic dysfunction (serum liver enzyme >3 times the normal limit)
- Renal failure (eGFR $<15 \text{ ml/min/1.73m}^2$ or requiring dialysis)
- ... and 4 more (see full listing online)

Locations (1 total)

Affiliated Hospital of Zunyi Medical University, Zunyi, Guizhou, China